



# Water Resources Data Wisconsin

## Water Year 1981



**U.S. GEOLOGICAL SURVEY WATER-DATA REPORT WI-81-1**  
Prepared in cooperation with the State of Wisconsin

## CALENDAR FOR WATER YEAR 1981

1980

### OCTOBER

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

### NOVEMBER

S	M	T	W	T	F	S
			2	3	4	5
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
	30					

### DECEMBER

S	M	T	W	T	F	S
			1		2	3
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			

1981

### JANUARY

S	M	T	W	T	F	S
			1	2	3	4
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

### FEBRUARY

S	M	T	W	T	F	S
			1	2	3	4
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28

### MARCH

S	M	T	W	T	F	S
			1	2	3	4
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

### APRIL

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30		

### MAY

S	M	T	W	T	F	S
			1	2		
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30

### JUNE

S	M	T	W	T	F	S
			1	2	3	4
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30				

### JULY

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

### AUGUST

S	M	T	W	T	F	S
			1			
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29

### SEPTEMBER

S	M	T	W	T	F	S
			1	2	3	4
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30			









































streams. Recorded low flows were generally above the 2-year frequency. An exception was the streams that flow into Lake Superior. These streams had low flows in the 2- to 5-year frequency during mid-September.

Streamflow tended to be in the normal range in fall and winter. Spring runoff was generally below normal due to the lack of snow, with dry conditions developing in May. Streamflow increased to above normal in June, especially in the northern part of the State, due to heavy June storms. Periodic storms during the summer maintained higher than normal flow for most of the summer.

Runoff for the year ranged from 80 to 144 percent of a long-term average (fig. 1). Monthly and annual flow for this year is compared to median discharges for a 30-year base period (fig. 2).




Figure 1. 1981 runoff as percent of long-term average runoff.




Figure 2. Comparison of discharge at representative gaging stations during 1981 water year with median discharge for 1916-80.





**ST. LAWRENCE RIVER BASIN RECORDS**




Figure 4. Major surface-water drainage basins and index of hydrologic records.































































































































































































































































































































































































































































































































































































































































































































































































































October 1, 1978

## FACTORS FOR CONVERTING INCH-POUND UNITS TO INTERNATIONAL SYSTEM UNITS (SI)

The following factors may be used to convert the inch-pound units published herein to the International System of Units (SI). This report contains both the inch-pound and SI unit equivalents in the station manuscript descriptions.

Multiply inch-pound units	By	To obtain SI units
<i>Length</i>		
inches (in)	$2.54 \times 10^1$	millimeters (mm)
	$2.54 \times 10^{-2}$	meters (m)
feet (ft)	$3.048 \times 10^{-1}$	meters (m)
miles (mi)	$1.609 \times 10^0$	kilometers (km)
<i>Area</i>		
acres	$4.047 \times 10^3$	square meters ( $m^2$ )
	$4.047 \times 10^{-1}$	square hectometers ( $hm^2$ )
	$4.047 \times 10^{-3}$	square kilometers ( $km^2$ )
square miles ( $mi^2$ )	$2.590 \times 10^0$	square kilometers ( $km^2$ )
<i>Volume</i>		
gallons (gal)	$3.785 \times 10^0$	liters (L)
	$3.785 \times 10^0$	cubic decimeters ( $dm^3$ )
	$3.785 \times 10^{-3}$	cubic meters ( $m^3$ )
million gallons	$3.785 \times 10^3$	cubic meters ( $m^3$ )
	$3.785 \times 10^{-3}$	cubic hectometers ( $hm^3$ )
cubic feet ( $ft^3$ )	$2.832 \times 10^1$	cubic decimeters ( $dm^3$ )
	$2.832 \times 10^{-2}$	cubic meters ( $m^3$ )
cfs-days	$2.447 \times 10^3$	cubic meters ( $m^3$ )
	$2.447 \times 10^{-3}$	cubic hectometers ( $hm^3$ )
acre-feet (acre-ft)	$1.233 \times 10^3$	cubic meters ( $m^3$ )
	$1.233 \times 10^{-3}$	cubic hectometers ( $hm^3$ )
	$1.233 \times 10^{-6}$	cubic kilometers ( $km^3$ )
<i>Flow</i>		
cubic feet per second ( $ft^3/s$ )	$2.832 \times 10^1$	liters per second (L/s)
	$2.832 \times 10^1$	cubic decimeters per second ( $dm^3/s$ )
	$2.832 \times 10^{-2}$	cubic meters per second ( $m^3/s$ )
gallons per minute (gal/min)	$6.309 \times 10^{-2}$	liters per second (L/s)
	$6.309 \times 10^{-2}$	cubic decimeters per second ( $dm^3/s$ )
	$6.309 \times 10^{-5}$	cubic meters per second ( $m^3/s$ )
million gallons per day	$4.381 \times 10^1$	cubic decimeters per second ( $dm^3/s$ )
	$4.381 \times 10^{-2}$	cubic meters per second ( $m^3/s$ )
<i>Mass</i>		
tons (short)	$9.072 \times 10^{-1}$	megagrams (Mg) or metric tons

POSTAGE AND FEES PAID  
U.S. DEPARTMENT OF THE INTERIOR  
INT 413



U.S. DEPARTMENT OF THE INTERIOR  
Geological Survey  
1815 University Avenue, Room 200  
Madison, WI 53706

OFFICIAL BUSINESS  
PENALTY FOR PRIVATE USE \$300  
SPECIAL 4TH CLASS BOOK RATE